

CLNE0026: Treatment (HASU and Service Delivery)

Old code : CLNEG056

View Online



[1]

'Stroke Series | ACNR | Online Neurology Journal'. [Online]. Available: <http://www.acnr.co.uk/category/stroke-series/>

[2]

P. Gates, 'The rule of 4 of the brainstem: a simplified method for understanding brainstem anatomy and brainstem vascular syndromes for the non-neurologist', *Internal Medicine Journal*, vol. 35, no. 4, pp. 263–266, Apr. 2005, doi: 10.1111/j.1445-5994.2004.00732.x.

[3]

J. Patten, *Neurological differential diagnosis*, 2nd ed. London: Springer, 1996.

[4]

H.J.M. Barnett, 'The classification of ischaemic stroke', in *Stroke: pathophysiology, diagnosis and management*, 3rd ed., New York ; Edinburgh: Churchill Livingstone, 1998.

[5]

C. Warlow et al., Eds, 'What Pathological Type of Stroke is it, Cerebral Ischaemia or Haemorrhage?', in *Stroke*, Oxford, UK: Blackwell Publishing Ltd, 2008, pp. 181–258. doi: 10.1002/9780470696361.ch5.

[6]

J. C. Hemphill et al., 'Guidelines for the Management of Spontaneous Intracerebral Hemorrhage', *Stroke*, vol. 46, no. 7, pp. 2032–2060, Jul. 2015, doi: 10.1161/STR.0000000000000069.

[7]

P. Bhogal, T. Andersson, V. Maus, A. Mpotsaris, and L. Yeo, 'Mechanical Thrombectomy—A Brief Review of a Revolutionary new Treatment for Thromboembolic Stroke', *Clinical Neuroradiology*, vol. 28, no. 3, pp. 313–326, Sep. 2018, doi: 10.1007/s00062-018-0692-2.

[8]

P. Bhogal et al., 'Mechanical thrombectomy in children and adolescents: report of five cases and literature review', *Stroke and Vascular Neurology*, Nov. 2018, doi: 10.1136/svn-2018-000181.

[9]

P. Bhogal et al., 'A Comparison of Mechanical Thrombectomy in the M1 and M2 Segments of the Middle Cerebral Artery: A Review of 585 Consecutive Patients', *Interventional Neurology*, vol. 6, no. 3–4, pp. 191–198, 2017, doi: 10.1159/000475535.

[10]

C. Foerch, 'Survival and quality of life outcome after mechanical ventilation in elderly stroke patients', *Journal of Neurology, Neurosurgery & Psychiatry*, vol. 75, no. 7, pp. 988–993, Jul. 2004, doi: 10.1136/jnnp.2003.021014.

[11]

J. R. Armstrong and B. D. Mosher, 'Aspiration Pneumonia After Stroke', *The Neurohospitalist*, vol. 1, no. 2, pp. 85–93, Apr. 2011, doi: 10.1177/1941875210395775.

[12]

P. J. Friedman, 'Atrial fibrillation after stroke in the elderly.', *Stroke*, vol. 22, no. 2, pp. 209–214, Feb. 1991, doi: 10.1161/01.STR.22.2.209.

[13]

'Gastrointestinal bleeding in stroke. - PubMed - NCBI'. [Online]. Available: https://ovidsp.tx.ovid.com/sp-3.32.0a/ovidweb.cgi?&S=PEHDFPHICLDDJPAINCDKPAOBNHOKAA00&Link+Set=S.sh.22.23.27.31.38.44%7c7%7csl_10

[14]

M. S. Ekker et al., 'Epidemiology, aetiology, and management of ischaemic stroke in young adults', *The Lancet Neurology*, vol. 17, no. 9, pp. 790–801, Sep. 2018, doi: 10.1016/S1474-4422(18)30233-3.

[15]

J. M. Ferro, A. R. Massaro, and J.-L. Mas, 'Aetiological diagnosis of ischaemic stroke in young adults', *The Lancet Neurology*, vol. 9, no. 11, pp. 1085–1096, Nov. 2010, doi: 10.1016/S1474-4422(10)70251-9.

[16]

J. Putaala et al., 'Analysis of 1008 Consecutive Patients Aged 15 to 49 With First-Ever Ischemic Stroke', *Stroke*, vol. 40, no. 4, pp. 1195–1203, Apr. 2009, doi: 10.1161/STROKEAHA.108.529883.

[17]

Y.-C. Cheng et al., 'Cocaine Use and Risk of Ischemic Stroke in Young Adults', *Stroke*, vol. 47, no. 4, pp. 918–922, Apr. 2016, doi: 10.1161/STROKEAHA.115.011417.

[18]

A. B. Mitchell et al., 'Obesity Increases Risk of Ischemic Stroke in Young Adults', *Stroke*, vol. 46, no. 6, pp. 1690–1692, Jun. 2015, doi: 10.1161/STROKEAHA.115.008940.

[19]

R. Roivainen, E. Haapaniemi, J. Putaala, M. Kaste, and T. Tatlisumak, 'Young adult

ischaemic stroke related acute symptomatic and late seizures: risk factors', *European Journal of Neurology*, vol. 20, no. 9, pp. 1247–1255, Sep. 2013, doi: 10.1111/ene.12157.

[20]

B. S. Jacobs, B. Boden-Albala, I.-F. Lin, and R. L. Sacco, 'Stroke in the Young in the Northern Manhattan Stroke Study', *Stroke*, vol. 33, no. 12, pp. 2789–2793, Dec. 2002, doi: 10.1161/01.STR.0000038988.64376.3A.

[21]

A. Towfighi, J. L. Saver, R. Engelhardt, and B. Ovbiagele, 'Factors Associated with the Steep Increase in Late Midlife Stroke Occurrence among US Men', *Journal of Stroke and Cerebrovascular Diseases*, vol. 17, no. 4, pp. 165–168, Jul. 2008, doi: 10.1016/j.jstrokecerebrovasdis.2007.12.007.

[22]

N. Allen, 'Racial/Ethnic Differences in Stroke in Young Adults', *Neuroepidemiology*, vol. 32, no. 4, pp. 312–312, 2009, doi: 10.1159/000208796.

[23]

M. Arnold et al., 'Age-dependent differences in demographics, risk factors, co-morbidity, etiology, management, and clinical outcome of acute ischemic stroke', *Journal of Neurology*, vol. 255, no. 10, pp. 1503–1507, Oct. 2008, doi: 10.1007/s00415-008-0949-9.

[24]

N. K. de Rooij, F. H. H. Linn, J. A. van der Plas, A. Algra, and G. J. E. Rinkel, 'Incidence of subarachnoid haemorrhage: a systematic review with emphasis on region, age, gender and time trends', *Journal of Neurology, Neurosurgery & Psychiatry*, vol. 78, no. 12, pp. 1365–1372, Dec. 2007, doi: 10.1136/jnnp.2007.117655.

[25]

By: Kissela, BM (Kissela, Brett M.)[1] ; Alwell, K (Alwell, Kathleen)[1] ;

Khoury, J (Khoury, Jane)[1] ; Moomaw, CJ (Moomaw, Charles J.)[1] ;
 Woo, D (Woo, Daniel)[1] ; Adeoye, O (Adeoye, Opeolu)[1] ; Flaherty,
 ML (Flaherty, Matthew L.)[1] ; Khatri, P (Khatri, Pooja)[1] ; Ferioli, S
 (Ferioli, Simona)[1] ; Broderick, JP (Broderick, Joseph P.)[1] ;
 Kleindorfer, D (Kleindorfer, Dawn)[1] ...More...Less, 'Stroke in the Young is
 Increasing Over Time: Temporal Trends in the Age of Stroke Incidence in a Large, Bi-racial
 Population', STROKE STROKE, vol. 41, no. 4, 2010, doi: 10.

[26]

A. Merwick and D. Werring, 'Posterior circulation ischaemic stroke', BMJ, vol. 348, no.
 may19 33, pp. g3175–g3175, May 2014, doi: 10.1136/bmj.g3175.

[27]

S. D. Treadwell, B. Thanvi, and T. G. Robinson, 'Stroke in pregnancy and the puerperium',
 Postgraduate Medical Journal, vol. 84, no. 991, pp. 238–245, May 2008, doi:
 10.1136/pgmj.2007.066167.

[28]

M. Wasay et al., 'Ischemic Stroke in Young Asian Women: Risk Factors, Subtypes and
 Outcome', Cerebrovascular Diseases, vol. 30, no. 4, pp. 418–422, 2010, doi:
 10.1159/000317075.

[29]

N. Pérez de la Ossa et al., 'Design and Validation of a Prehospital Stroke Scale to Predict
 Large Arterial Occlusion', Stroke, vol. 45, no. 1, pp. 87–91, Jan. 2014, doi:
 10.1161/STROKEAHA.113.003071.

[30]

M. Goyal et al., 'Endovascular thrombectomy after large-vessel ischaemic stroke: a
 meta-analysis of individual patient data from five randomised trials', The Lancet, vol. 387,
 no. 10029, pp. 1723–1731, Apr. 2016, doi: 10.1016/S0140-6736(16)00163-X.

[31]

J. L. Saver et al., 'Time to Treatment With Endovascular Thrombectomy and Outcomes From Ischemic Stroke: A Meta-analysis', JAMA, vol. 316, no. 12, Sep. 2016, doi: 10.1001/jama.2016.13647.

[32]

J. Stone, 'Functional neurological disorders: the neurological assessment as treatment', Practical Neurology, vol. 16, no. 1, pp. 7–17, Feb. 2016, doi: 10.1136/practneurol-2015-001241.

[33]

A. Carson, A. Lehn, L. Ludwig, and J. Stone, 'Explaining functional disorders in the neurology clinic: a photo story', Practical Neurology, vol. 16, no. 1, pp. 56–61, Feb. 2016, doi: 10.1136/practneurol-2015-001242.

[34]

M. J. Edwards and K. P. Bhatia, 'Functional (psychogenic) movement disorders: merging mind and brain', The Lancet Neurology, vol. 11, no. 3, pp. 250–260, Mar. 2012, doi: 10.1016/S1474-4422(11)70310-6.

[35]

M. J. Edwards, R. A. Adams, H. Brown, I. Parees, and K. J. Friston, 'A Bayesian account of "hysteria"', Brain, vol. 135, no. 11, pp. 3495–3512, Nov. 2012, doi: 10.1093/brain/aws129.

[36]

J. Stone, 'Functional weakness and sensory disturbance', Journal of Neurology, Neurosurgery & Psychiatry, vol. 73, no. 3, pp. 241–245, Sep. 2002, doi: 10.1136/jnnp.73.3.241.

[37]

G. Habib et al., '2015 ESC Guidelines for the management of infective endocarditis', European Heart Journal, vol. 36, no. 44, pp. 3075–3128, Nov. 2015, doi: 10.1093/eurheartj/ehv319.

[38]

J. M. Ferro and P. Canhão, 'Cerebral Venous Sinus Thrombosis: Update on Diagnosis and Management', *Current Cardiology Reports*, vol. 16, no. 9, Sep. 2014, doi: 10.1007/s11886-014-0523-2.

[39]

C. E. Robertson, R. D. Brown, E. F. M. Wijdicks, and A. A. Rabinstein, 'Recovery after spinal cord infarcts: Long-term outcome in 115 patients', *Neurology*, vol. 78, no. 2, pp. 114–121, Jan. 2012, doi: 10.1212/WNL.0b013e31823efc93.

[40]

D. G. Smithard, N. C. Smeeton, and C. D. A. Wolfe, 'Long-term outcome after stroke: does dysphagia matter?', *Age and Ageing*, vol. 36, no. 1, pp. 90–94, Nov. 2006, doi: 10.1093/ageing/afl149.

[41]

O. Ekberg, S. Hamdy, V. Woisard, A. Wuttge-Hannig, and P. Ortega, 'Social and Psychological Burden of Dysphagia: Its Impact on Diagnosis and Treatment', *Dysphagia*, vol. 17, no. 2, pp. 139–146, Apr. 2002, doi: 10.1007/s00455-001-0113-5.

[42]

S. HAMDY, J. C. ROTHWELL, Q. AZIZ, and D. G. THOMPSON, 'Organization and reorganization of human swallowing motor cortex: implications for recovery after stroke*', *Clinical Science*, vol. 99, no. 2, pp. 151–157, Aug. 2000, doi: 10.1042/cs0990151.

[43]

V. Steinhagen, A. Grossmann, R. Benecke, and U. Walter, 'Swallowing Disturbance Pattern Relates to Brain Lesion Location in Acute Stroke Patients', *Stroke*, vol. 40, no. 5, pp. 1903–1906, May 2009, doi: 10.1161/STROKEAHA.108.535468.

[44]

J. A. Hinchey, T. Shephard, K. Furie, D. Smith, D. Wang, and S. Tonn, 'Formal Dysphagia Screening Protocols Prevent Pneumonia', *Stroke*, vol. 36, no. 9, pp. 1972–1976, 2005, [Online]. Available: https://www.ahajournals.org/doi/full/10.1161/01.STR.0000177529.86868.8d?url_ver=Z39.88-2003&rfr_id=ori%3Arid%3Acrossref.org&rfr_dat=cr_pub%3Dpubmed

[45]

S. B. Leder, 'Videofluoroscopic Evaluation of Aspiration with Visual Examination of the Gag Reflex and Velar Movement', *Dysphagia*, vol. 12, no. 1, pp. 21–23, Jan. 1997, doi: 10.1007/PL00009514.

[46]

T. Warnecke et al., 'The Safety of Fiberoptic Endoscopic Evaluation of Swallowing in Acute Stroke Patients', *Stroke*, vol. 40, no. 2, pp. 482–486, Feb. 2009, doi: 10.1161/STROKEAHA.108.520775.

[47]

Reilly, S., Douglas, J., and Oates J., 'The evidence base for the management of dysphagia', in *Evidence-based practice in speech pathology*, London: Whurr, 2004. [Online]. Available: <https://contentstore.cla.co.uk/secure/link?id=c19574b4-dd02-e911-80cd-005056af4099>

[48]

S. Broadley et al., 'Predictors of prolonged dysphagia following acute stroke', *Journal of Clinical Neuroscience*, vol. 10, no. 3, pp. 300–305, May 2003, doi: 10.1016/S0967-5868(03)00022-5.

[49]

K. Toyoda, M. Koga, and S. Sato, 'What are the next breakthroughs in the management of acute intracerebral hemorrhage?', *International Journal of Stroke*, vol. 11, no. 4, pp. 386–388, Jun. 2016, doi: 10.1177/1747493016631559.

[50]

A. D. Mendelow, B. A. Gregson, E. N. Rowan, G. D. Murray, A. Gholkar, and P. M. Mitchell, 'Early surgery versus initial conservative treatment in patients with spontaneous supratentorial lobar intracerebral haematomas (STICH II): a randomised trial', *The Lancet*, vol. 382, no. 9890, pp. 397–408, Aug. 2013, doi: 10.1016/S0140-6736(13)60986-1.

[51]

STICH investigatorsMendelow, A DavidGregson, Barbara AFernandes, Helen MMurray, Gordon D, 'Early surgery versus initial conservative treatment in patients with spontaneous supratentorial intracerebral haematomas in the International Surgical Trial in Intracerebral Haemorrhage (STICH): a randomised trial', *The Lancet*, vol. 365, no. 5, pp. 387–97, [Online]. Available: https://search.proquest.com/docview/198997243?rfr_id=info%3Axri%2Fsid%3Aprim

[52]

D. Fiorella, A. Arthur, M. Bain, and J. Mocco, 'Minimally Invasive Surgery for Intracerebral and Intraventricular Hemorrhage', *Stroke*, vol. 47, no. 5, pp. 1399–1406, May 2016, doi: 10.1161/STROKEAHA.115.011415.

[53]

C. Bushnell et al., 'Guidelines for the Prevention of Stroke in Women', *Stroke*, vol. 45, no. 5, pp. 1545–1588, May 2014, doi: 10.1161/01.str.0000442009.06663.48.

[54]

S. D. Treadwell, B. Thanvi, and T. G. Robinson, 'Stroke in pregnancy and the puerperium', *Postgraduate Medical Journal*, vol. 84, no. 991, pp. 238–245, May 2008, doi: 10.1136/pgmj.2007.066167.

[55]

Z. Moatti, M. Gupta, R. Yadava, and S. Thamban, 'A review of stroke and pregnancy: incidence, management and prevention', *European Journal of Obstetrics & Gynecology and Reproductive Biology*, vol. 181, pp. 20–27, Oct. 2014, doi: 10.1016/j.ejogrb.2014.07.024.

[56]

E. Del Zotto, A. Giossi, I. Volonghi, P. Costa, A. Padovani, and A. Pezzini, 'Ischemic Stroke during Pregnancy and Puerperium', *Stroke Research and Treatment*, vol. 2011, pp. 1–13, 2011, doi: 10.4061/2011/606780.

[57]

M. A. O'Neal and S. K. Feske, 'Stroke in pregnancy: a case-oriented review', *Practical Neurology*, vol. 16, no. 1, pp. 23–34, Feb. 2016, doi: 10.1136/practneurol-2015-001217.

[58]

M. E. van Alebeek, R. de Heus, A. M. Tuladhar, and F.-E. de Leeuw, 'Pregnancy and ischemic stroke', *Current Opinion in Neurology*, Nov. 2017, doi: 10.1097/WCO.0000000000000522.

[59]

R. H. Swartz et al., 'Canadian stroke best practice consensus statement: Secondary stroke prevention during pregnancy', *International Journal of Stroke*, vol. 13, no. 4, pp. 406–419, Jun. 2018, doi: 10.1177/1747493017743801.

[60]

'Stroke rehabilitation in adults | Guidance and guidelines | NICE', [Online]. Available: <https://www.nice.org.uk/guidance/cg162>

[61]

P. Langhorne and S. Baylan, 'Early supported discharge services for people with acute stroke', *Cochrane Database of Systematic Reviews*, Jul. 2017, doi: 10.1002/14651858.CD000443.pub4.

[62]

R. J. Fisher et al., 'Is Stroke Early Supported Discharge still effective in practice? A prospective comparative study', *Clinical Rehabilitation*, vol. 30, no. 3, pp. 268–276, Mar. 2016, doi: 10.1177/0269215515578697.

[63]

'Reimagining community services | The King's Fund'. [Online]. Available:
<https://www.kingsfund.org.uk/publications/community-services-assets>

[64]

'The key principles of effective discharge planning | Clinical | Nursing Times'. [Online]. Available:
<https://www.nursingtimes.net/roles/nurse-managers/the-key-principles-of-effective-discharge-planning/5053740.article>

[65]

'Transition between inpatient hospital settings and community or care home settings for adults with social care needs | Guidance and guidelines | NICE', [Online]. Available:
<https://www.nice.org.uk/guidance/ng27>

[66]

'Organised inpatient (stroke unit) care for stroke', Cochrane Database of Systematic Reviews, Sep. 2013, doi: 10.1002/14651858.CD000197.pub3.

[67]

B. D. Bray et al., 'Associations between the organisation of stroke services, process of care, and mortality in England: prospective cohort study', BMJ, vol. 346, no. may10 1, pp. f2827-f2827, May 2013, doi: 10.1136/bmj.f2827.

[68]

R. M. Hunter et al., 'Impact on Clinical and Cost Outcomes of a Centralized Approach to Acute Stroke Care in London: A Comparative Effectiveness Before and After Model', PLoS ONE, vol. 8, no. 8, Aug. 2013, doi: 10.1371/journal.pone.0070420.

[69]

S. Morris et al., 'Impact of centralising acute stroke services in English metropolitan areas on mortality and length of hospital stay: difference-in-differences analysis', *BMJ*, vol. 349, no. aug04 4, pp. g4757–g4757, Aug. 2014, doi: 10.1136/bmj.g4757.

[70]

'Neuropsychological assessment.', *Journal of Neurology, Neurosurgery, and Psychiatry*, vol. 58, no. 6, 1995, [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1073539/>

[71]

P. S. Sachdev, D. M. Lipnicki, J. D. Crawford, W. Wen, and H. Brodaty, 'Progression of cognitive impairment in stroke/TIA patients over 3 years', *Journal of Neurology, Neurosurgery & Psychiatry*, vol. 85, no. 12, pp. 1324–1330, Dec. 2014, doi: 10.1136/jnnp-2013-306776.

[72]

G. M. S. Nys et al., 'The prognostic value of domain-specific cognitive abilities in acute first-ever stroke', *Neurology*, vol. 64, no. 5, pp. 821–827, Mar. 2005, doi: 10.1212/01.WNL.0000152984.28420.5A.

[73]

R. Hurford, A. Charidimou, Z. Fox, L. Cipolotti, and D. J. Werring, 'Domain-specific trends in cognitive impairment after acute ischaemic stroke', *Journal of Neurology*, vol. 260, no. 1, pp. 237–241, Jan. 2013, doi: 10.1007/s00415-012-6625-0.

[74]

C. M. Van Heugten, L. Walton, and U. Hentschel, 'Can we forget the Mini-Mental State Examination? A systematic review of the validity of cognitive screening instruments within one month after stroke', *Clinical Rehabilitation*, vol. 29, no. 7, pp. 694–704, Jul. 2015, doi: 10.1177/0269215514553012.

[75]

L. Burton and S. Tyson, 'Screening for cognitive impairment after stroke: A systematic

review of psychometric properties and clinical utility', *Journal of Rehabilitation Medicine*, vol. 47, no. 3, pp. 193–203, 2015, doi: 10.2340/16501977-1930.

[76]

E. Chan, S. Altendorff, C. Healy, D. J. Werring, and L. Cipolotti, 'The test accuracy of the Montreal Cognitive Assessment (MoCA) by stroke lateralisation', *Journal of the Neurological Sciences*, vol. 373, pp. 100–104, Feb. 2017, doi: 10.1016/j.jns.2016.12.028.

[77]

E. Chan, S. Khan, R. Oliver, S. K. Gill, D. J. Werring, and L. Cipolotti, 'Underestimation of cognitive impairments by the Montreal Cognitive Assessment (MoCA) in an acute stroke unit population', *Journal of the Neurological Sciences*, vol. 343, no. 1–2, pp. 176–179, Aug. 2014, doi: 10.1016/j.jns.2014.05.005.

[78]

F. C. Ng et al., 'Deconstruction of Interhospital Transfer Workflow in Large Vessel Occlusion', *Stroke*, vol. 48, no. 7, pp. 1976–1979, Jul. 2017, doi: 10.1161/STROKEAHA.117.017235.

[79]

N. Kamal, E. E. Smith, T. Jeerakathil, and M. D. Hill, 'Thrombolysis: Improving door-to-needle times for ischemic stroke treatment – A narrative review', *International Journal of Stroke*, vol. 13, no. 3, pp. 268–276, Apr. 2018, doi: 10.1177/1747493017743060.

[80]

N. Kamal et al., 'Good is not Good Enough: The Benchmark Stroke Door-to-Needle Time Should be 30 Minutes', *Canadian Journal of Neurological Sciences / Journal Canadien des Sciences Neurologiques*, vol. 41, no. 06, pp. 694–696, Nov. 2014, doi: 10.1017/cjn.2014.41.

[81]

K. Salter, C. Hellings, N. Foley, and R. Teasell, 'The experience of living with stroke: a

qualitative meta-synthesis', *Journal of Rehabilitation Medicine*, vol. 40, no. 8, pp. 595–602, 2008, doi: 10.2340/16501977-0238.

[82]

S. Simeone, S. Savini, M. Z. Cohen, R. Alvaro, and E. Vellone, 'The experience of stroke survivors three months after being discharged home: A phenomenological investigation', *European Journal of Cardiovascular Nursing*, vol. 14, no. 2, pp. 162–169, Apr. 2015, doi: 10.1177/1474515114522886.

[83]

B. Norrving et al., 'Cross-National Key Performance Measures of the Quality of Acute Stroke Care in Western Europe', *Stroke*, vol. 46, no. 10, pp. 2891–2895, Oct. 2015, doi: 10.1161/STROKEAHA.115.008811.

[84]

'Quality improvement made simple | The Health Foundation'. [Online]. Available: <https://www.health.org.uk/publications/quality-improvement-made-simple>

[85]

'Mind the Gap! Third SSNAP Annual Report'. [Online]. Available: https://kingsfund.blogs.com/health_management/2016/11/mind-the-gap-the-third-ssnap-annual-report-care-received-between-april-2015-to-march-2016.html

[86]

P. Lindsay, K. L. Furie, S. M. Davis, G. A. Donnan, and B. Norrving, 'World Stroke Organization Global Stroke Services Guidelines and Action Plan', *International Journal of Stroke*, vol. 9, no. SA100, pp. 4–13, Oct. 2014, doi: 10.1111/ijss.12371.

[87]

B. D. Bray et al., 'Associations between the organisation of stroke services, process of care, and mortality in England: prospective cohort study', *BMJ*, vol. 346, no. may10 1, pp. f2827–f2827, May 2013, doi: 10.1136/bmj.f2827.

[88]

'Sentinel Stroke National Audit Programme (SSNAP)'. [Online]. Available:
<https://www.strokeaudit.org/>

[89]

'Scottish Stroke Care Audit (SSCA)'. [Online]. Available:
<https://www.strokeaudit.scot.nhs.uk/index.html>