

PSYCGN42: Evaluating Clinical Interventions. Lecturer: Julian Edbrooke-Childs

View Online



[1]

M. Wolpert et al., 'Drawing on the evidence: Advice for mental health professionals working with children and adolescents'. CAMHS Publications, London, 2006. Available: https://www.ucl.ac.uk/ebpu/docs/publication_files/drawing_onthe_evidence

[2]

B. F. Chorpita et al., 'Evidence-based treatments for children and adolescents: An updated review of indicators of efficacy and effectiveness', *Clinical Psychology: Science and Practice*, vol. 18, no. 2, pp. 154-172, 2011, doi: 10.1111/j.1468-2850.2011.01247.x

[3]

A. E. Kazdin, 'Evidence-based treatment and practice: New opportunities to bridge clinical research and practice, enhance the knowledge base, and improve patient care.', *American Psychologist*, vol. 63, no. 3, pp. 146-159, 2008, Available: <http://ovidsp.ovid.com.libproxy.ucl.ac.uk/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&AN=00000487-200804000-00001&LSLINK=80&D=ovft>

[4]

A. Girard, A. Rochette, and B. Fillion, 'Knowledge translation and improving practices in neurological rehabilitation: managers' viewpoint', *Journal of Evaluation in Clinical Practice*, vol. 19, no. 1, pp. 60-67, 2013, doi: 10.1111/j.1365-2753.2011.01769.x

[5]

K. P. Gallo and D. H. Barlow, 'Factors involved in clinician adoption and nonadoption of

evidence-based interventions in mental health', *Clinical Psychology: Science and Practice*, vol. 19, no. 1, pp. 93–106, 2012, doi: 10.1111/j.1468-2850.2012.01276.x

[6]

L. Timulak, 'Randomized control trials', in *Research in psychotherapy and counselling*, Los Angeles: SAGE, 2008, pp. 30–43. Available:
<https://contentstore.cla.co.uk/secure/link?id=3b67bef9-7b36-e711-80c9-005056af4099>

[7]

M. Cooper, 'Meeting the demand for evidence-based practice', *Therapy Today*, vol. 22, no. 4, pp. 10–16, Available: <http://strathprints.strath.ac.uk/32220/>

[8]

Wolpert, M., Fugard, A. J. B., and Deighton, J., 'Issues in evaluation of psychotherapies', in *Cognitive behaviour therapy for children and families*, P. J. Graham and S. Reynolds, Eds, 3rd ed. Cambridge: Cambridge University Press, 2013, pp. 34–47. Available:
<https://contentstore.cla.co.uk/secure/link?id=0ccb9e2-7836-e711-80c9-005056af4099>

[9]

J. R. Weisz et al., 'Performance of evidence-based youth psychotherapies compared with usual clinical care', *JAMA Psychiatry*, vol. 70, no. 7, 2013, doi:
10.1001/jamapsychiatry.2013.1176

[10]

A. Liberati et al., 'The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration', *BMJ*, vol. 339, no. jul21 1, pp. b2700–b2700, 2009, doi: 10.1136/bmj.b2700

[11]

Improving Access to Psychological Therapies (IAPT) and Department of Health, 'IAPT 3 Year Report - The first million patients. November 2012'. Department of Health, 2012. Available:
<http://webarchive.nationalarchives.gov.uk/20160302154833/http://www.iapt.nhs.uk/silo/fil>

es/iapt-3-year-report.pdf

[12]

J. Bullivant et al., 'Clinical audit: A guide for NHS Boards and partners.' : Healthcare Quality Improvement Partnership and Good Governance Institute., London, 2015. Available: <https://www.hqip.org.uk/resources/clinical-audit-a-guide-for-nhs-boards-and-partners/>

[13]

S.-E. Kam and N. Midgley, 'Exploring clinical judgement: How do child and adolescent mental health professionals decide whether a young person needs individual psychotherapy?', *Clinical Child Psychology and Psychiatry*, vol. 11, no. 1, pp. 27–44, 2006, doi: 10.1177/1359104506059122

[14]

M. Semmelroch et al, 'Retrospective audit of postictal generalized EEG suppression in telemetry', *Epilepsia*, vol. 53, no. 2, pp. e21–e24, 2012, doi: 10.1111/j.1528-1167.2011.03296.x

[15]

J. Edbrooke-Childs, M. Wolpert, and J. Deighton, 'Using Patient Reported Outcome Measures to Improve Service Effectiveness (UPROMISE): Training clinicians to use outcome measures in child mental health', *Administration and Policy in Mental Health and Mental Health Services Research*, vol. Online publication, Oct. 2014, doi: 10.1007/s10488-014-0600-2

[16]

L. Bickman et al, 'Effects of routine feedback to clinicians on mental health outcomes of youths: Results of a randomized trial', *Psychiatric Services*, vol. 62, no. 12, pp. 1423–1429, 2011, doi: 10.1176/appi.ps.002052011

[17]

K. de Jong et al., 'Understanding the differential impact of outcome monitoring: Therapist variables that moderate feedback effects in a randomized clinical trial', *Psychotherapy*

Research, vol. 22, no. 4, pp. 464–474, 2012, doi: 10.1080/10503307.2012.673023

[18]

P. Moran et al, 'What do parents and carers think about routine outcome measures and their use? A focus group study of CAMHS attenders', *Clinical Child Psychology and Psychiatry*, vol. 17, no. 1, pp. 65–79, 2012, doi: 10.1177/1359104510391859

[19]

M. Wolpert, K. Curtis-Tyler, and J. Edbrooke-Childs, 'A qualitative exploration of patient and clinician views on Patient Reported Outcome Measures in child mental health and diabetes services', *Administration and Policy in Mental Health and Mental Health Services Research*, vol. Online publication, Sept. 2014, doi: 10.1007/s10488-014-0586-9

[20]

D. Scheinost et al., 'Orbitofrontal cortex neurofeedback produces lasting changes in contamination anxiety and resting-state connectivity', *Translational Psychiatry*, vol. 3, no. 4, 2013, doi: 10.1038/tp.2013.24

[21]

Young Minds, 'Talking about talking therapies: Thinking and planning about how to make good and accessible talking therapies available to children and young people.' Available: <http://www.iapt.nhs.uk/silo/files/talking-about-talking-therapies.pdf>

[22]

C. Evans, F. Margison, and M. Barkham, 'The contribution of reliable and clinically significant change methods to evidence-based mental health', *Evidence-Based Mental Health*, vol. 1, no. 3, pp. 70–72, 1998, doi: 10.1136/ebmh.1.3.70

[23]

T. Ford et al, 'Strengths and Difficulties Questionnaire Added Value Scores: Evaluating effectiveness in child mental health interventions', *The British Journal of Psychiatry*, vol. 194, no. 6, pp. 552–558, 2009, doi: 10.1192/bjp.bp.108.052373

[24]

Child Outcomes Research Consortium (CORC), 'CORC+ Annual report: Erinsborough'. CORC, 2014. Available: <http://www.corc.uk.net/resources/downloads/>

[25]

M. J. Lambert and B. M. Ogles, 'Using clinical significance in psychotherapy outcome research: The need for a common procedure and validity data', *Psychotherapy Research*, vol. 19, no. 4-5, pp. 493-501, 2009, doi: 10.1080/10503300902849483

[26]

'CAMHS Outcomes Research Consortium | Collaborating to improve quality through outcomes'. Available: <http://www.corc.uk.net/>

[27]

I. Fleming et al., 'Learning from a learning collaboration: The CORC approach to combining research, evaluation and practice in child mental health', *Administration and Policy in Mental Health and Mental Health Services Research*, vol. Online publication, Sept. 2014, doi: 10.1007/s10488-014-0592-y

[28]

M. J. Batty et al, 'Implementing routine outcome measures in child and adolescent mental health services: from present to future practice', *Child and Adolescent Mental Health*, vol. 18, no. 2, pp. 82-87, 2013, doi: 10.1111/j.1475-3588.2012.00658.x

[29]

M. Jones and et al, 'Current view tool: Completion guide'. CAMHS Press, London, 2013. Available: <https://www.ucl.ac.uk/ebpu/publications/professionals>

[30]

J. Mellor-Clark et al., 'Leading horses to water: Lessons from a decade of helping

psychological therapy services use routine outcome measurement to improve practice', Administration and Policy in Mental Health and Mental Health Services Research, vol. Online publication, Sept. 2014, doi: 10.1007/s10488-014-0587-8

[31]

M. Rose and A. Bezjak, 'Logistics of collecting patient-reported outcomes (PROs) in clinical practice: an overview and practical examples', Quality of Life Research, vol. 18, no. 1, pp. 125-136, 2009, doi: 10.1007/s11136-008-9436-0

[32]

M. Wolpert et al, 'Routine outcomes monitoring as part of children and young people's Improving Access to Psychological Therapies (CYP IAPT) - improving care or unhelpful burden?', Child and Adolescent Mental Health, vol. 17, no. 3, pp. 129-130, 2012, doi: 10.1111/j.1475-3588.2012.00676.x

[33]

N. Abrines-Jaume et al., 'A qualitative analysis of implementing shared decision making in Child and Adolescent Mental Health Services in the United Kingdom: Stages and facilitators', Clinical Child Psychology and Psychiatry, vol. Advance online publication, Sept. 2014, doi: 10.1177/1359104514547596

[34]

O. Lindhiem et al., 'Client preferences affect treatment satisfaction, completion, and clinical outcome: A meta-analysis', Clinical Psychology Review, vol. 34, no. 6, pp. 506-517, 2014, doi: 10.1016/j.cpr.2014.06.002

[35]

D. Stacey et al., 'Decision aids for people facing health treatment or screening decisions', Cochrane Database of Systematic Reviews, no. 1, Art. CD001431, 2014, doi: 10.1002/14651858.CD001431.pub4. Available: <http://doi.wiley.com/10.1002/14651858.CD001431.pub4>

[36]

P. L. Wachtel, 'Beyond "ESTs": Problematic assumptions in the pursuit of evidence-based practice', *Psychoanalytic Psychology*, vol. 27, pp. 251–272, 2010.

[37]

N. Ivers et al., 'Audit and feedback: Effects on professional practice and healthcare outcomes', *Cochrane Database of Systematic Reviews*, no. 6, Art. CD000259, 2012, doi: 10.1002/14651858.CD000259.pub3. Available: <http://doi.wiley.com/10.1002/14651858.CD000259.pub3>