

CLNEG058: Neurorehabilitation

[View Online](#)

[1]

T. H. Murphy and D. Corbett, 'Plasticity during stroke recovery: from synapse to behaviour', *Nature Reviews Neuroscience*, vol. 10, no. 12, pp. 861–872, Dec. 2009, doi: 10.1038/nrn2735.

[2]

N. S. Ward, 'Restoring brain function after stroke — bridging the gap between animals and humans', *Nature Reviews Neurology*, vol. 13, no. 4, pp. 244–255, Apr. 2017, doi: 10.1038/nrneurol.2017.34.

[3]

D. J. Reinkensmeyer et al., 'Computational neurorehabilitation: modeling plasticity and learning to predict recovery', *Journal of NeuroEngineering and Rehabilitation*, vol. 13, no. 1, Dec. 2016, doi: 10.1186/s12984-016-0148-3.

[4]

Steven R Zeiler, 'The interaction between training and plasticity in the post-stroke brain', *Current opinion in neurology*, vol. 26, no. 6, 2013, doi: 10.1097/WCO.000000000000025. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4012223/>

[5]

T. Kitago and J. W. Krakauer, 'Motor learning principles for neurorehabilitation', in *Neurological Rehabilitation*, vol. 110, Elsevier, 2013, pp. 93–103 [Online]. Available: <http://linkinghub.elsevier.com/retrieve/pii/B9780444529015000083>

[6]

S. T. Carmichael, 'Brain Excitability in Stroke', *Archives of Neurology*, vol. 69, no. 2, Feb. 2012, doi: 10.1001/archneurol.2011.1175.

[7]

N. S. Ward, 'Does neuroimaging help to deliver better recovery of movement after stroke?', *Current Opinion in Neurology*, vol. 28, no. 4, pp. 323–329, Aug. 2015, doi: 10.1097/WCO.0000000000000223.

[8]

Archy O. de Berker, 'Predicting the behavioral impact of transcranial direct current stimulation: issues and limitations', *Frontiers in Human Neuroscience*, vol. 7, 2013, doi: 10.3389/fnhum.2013.00613. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3790257/>

[9]

S. K. Bhogal, R. Teasell, M. Speechley, and M. L. Albert, 'Intensity of Aphasia Therapy, Impact on Recovery * Aphasia Therapy Works!', *Stroke*, vol. 34, no. 4, pp. 987–993, Apr. 2003, doi: 10.1161/01.STR.0000062343.64383.D0.

[10]

M. C. Brady, H. Kelly, J. Godwin, P. Enderby, and P. Campbell, 'Speech and language therapy for aphasia following stroke', *Cochrane Database of Systematic Reviews*, Jun. 2016, doi: 10.1002/14651858.CD000425.pub4.

[11]

C. Breitenstein et al., 'Intensive speech and language therapy in patients with chronic aphasia after stroke: a randomised, open-label, blinded-endpoint, controlled trial in a health-care setting', *The Lancet*, vol. 389, no. 10078, pp. 1528–1538, Apr. 2017, doi: 10.1016/S0140-6736(17)30067-3.

[12]

J. Dignam et al., 'Intensive Versus Distributed Aphasia Therapy', *Stroke*, vol. 46, no. 8, pp.

2206–2211, Aug. 2015, doi: 10.1161/STROKEAHA.115.009522.

[13]

M. L. Berthier, 'Cognitive enhancing drugs in aphasia: A vote for hope', *Aphasiology*, vol. 28, no. 2, pp. 128–132, Feb. 2014, doi: 10.1080/02687038.2013.857756.

[14]

S. K. Bhogal, R. Teasell, M. Speechley, and M. L. Albert, 'Intensity of Aphasia Therapy, Impact on Recovery * Aphasia Therapy Works!', *Stroke*, vol. 34, no. 4, pp. 987–993, Apr. 2003, doi: 10.1161/01.STR.0000062343.64383.D0.

[15]

M. C. Brady, H. Kelly, J. Godwin, P. Enderby, and P. Campbell, 'Speech and language therapy for aphasia following stroke', *Cochrane Database of Systematic Reviews*, Jun. 2016, doi: 10.1002/14651858.CD000425.pub4.

[16]

P. S. Klonoff, *Psychotherapy after brain injury: principles and techniques*. New York: The Guilford Press, 2010.

[17]

'Quality Assurance Standards for physiotherapy service delivery | The Chartered Society of Physiotherapy'. [Online]. Available: <http://www.csp.org.uk/publications/quality-assurance-standards>

[18]

'Code of ethics and professional conduct'. [Online]. Available: https://www.rcot.co.uk/sites/default/files/CODE-OF-ETHICS-2015_0.pdf

[19]

S. E. F. James M., 'Contractures in orthopaedic and neurological conditions: a review of

causes and treatment', *Disability and Rehabilitation*, vol. 23, no. 13, pp. 549–558, Jan. 2001, doi: 10.1080/09638280010029930.

[20]

L. Harvey, I. de Jong, G. Goehl, and S. Marwedel, 'Twelve weeks of nightly stretch does not reduce thumb web-space contractures in people with a neurological condition: a randomised controlled trial', *Australian Journal of Physiotherapy*, vol. 52, no. 4, pp. 251–258, 2006, doi: 10.1016/S0004-9514(06)70004-6.

[21]

C. Kilbride, K. Hoffman, T. Baird, J. Tuckey, L. Marston, and L. D. Souza, 'Contemporary splinting practice in the UK for adults with neurological dysfunction: A cross-sectional survey', *International Journal of Therapy and Rehabilitation*, vol. 20, no. 11, pp. 559–566, Nov. 2013, doi: 10.12968/ijtr.2013.20.11.559.

[22]

N. A. Lannin, S. A. Horsley, R. Herbert, A. McCluskey, and A. Cusick, 'Splinting the hand in the functional position after brain impairment: A randomized, controlled trial', *Archives of Physical Medicine and Rehabilitation*, vol. 84, no. 2, pp. 297–302, Feb. 2003, doi: 10.1053/apmr.2003.50031.

[23]

N. A. Lannin, A. Cusick, A. McCluskey, and R. D. Herbert, 'Effects of Splinting on Wrist Contracture After Stroke: A Randomized Controlled Trial', *Stroke*, vol. 38, no. 1, pp. 111–116, Jan. 2007, doi: 10.1161/01.STR.0000251722.77088.12.

[24]

PA Mortenson, 'The use of casts in the management of joint mobility and hypertonia following brain injury in adults: a systematic review', 2003 [Online]. Available: <https://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0020153/>

[25]

P. Moorhouse and K. Rockwood, 'Vascular cognitive impairment: current concepts and clinical developments', *The Lancet Neurology*, vol. 7, no. 3, pp. 246–255, Mar. 2008, doi:

10.1016/S1474-4422(08)70040-1.

[26]

'Rehabilitation of cognitive impairment post stroke'. [Online]. Available: http://www.ebrsr.com/sites/default/files/Chapter%205_Cognitive.pdf

[27]

A. Chaudhuri and P. O. Behan, 'Fatigue in neurological disorders', *The Lancet*, vol. 363, no. 9413, pp. 978-988, Mar. 2004, doi: 10.1016/S0140-6736(04)15794-2.

[28]

F. Duncan, S. Wu, and G. E. Mead, 'Frequency and natural history of fatigue after stroke: A systematic review of longitudinal studies', *Journal of Psychosomatic Research*, vol. 73, no. 1, pp. 18-27, Jul. 2012, doi: 10.1016/j.jpsychores.2012.04.001.

[29]

W. De Doncker, R. Dantzer, H. Ormstad, and A. Kuppuswamy, 'Mechanisms of poststroke fatigue', *Journal of Neurology, Neurosurgery & Psychiatry*, Sep. 2017, doi: 10.1136/jnnp-2017-316007.

[30]

A. Kuppuswamy, 'The fatigue conundrum', *Brain*, vol. 140, no. 8, pp. 2240-2245, Aug. 2017, doi: 10.1093/brain/awx153.

[31]

A. Kuppuswamy, E. V. Clark, I. F. Turner, J. C. Rothwell, and N. S. Ward, 'Post-stroke fatigue: a deficit in corticomotor excitability?', *Brain*, vol. 138, no. 1, pp. 136-148, Jan. 2015, doi: 10.1093/brain/awu306.

[32]

A. Kuppuswamy, E. Clark, J. Rothwell, and N. S. Ward, 'Limb Heaviness',

Neurorehabilitation and Neural Repair, vol. 30, no. 4, pp. 360–362, May 2016, doi: 10.1177/1545968315597071.

[33]

'Clinical Guidelines for Stroke Management 2017'. [Online]. Available: <https://informme.org.au/Guidelines/Clinical-Guidelines-for-Stroke-Management-2017>

[34]

O. A. Howlett, N. A. Lannin, L. Ada, and C. McKinstry, 'Functional Electrical Stimulation Improves Activity After Stroke: A Systematic Review With Meta-Analysis', *Archives of Physical Medicine and Rehabilitation*, vol. 96, no. 5, pp. 934–943, May 2015, doi: 10.1016/j.apmr.2015.01.013.

[35]

J.-H. Lee, L. L. Baker, R. E. Johnson, and J. K. Tilson, 'Effectiveness of neuromuscular electrical stimulation for management of shoulder subluxation post-stroke: a systematic review with meta-analysis', *Clinical Rehabilitation*, vol. 31, no. 11, pp. 1431–1444, Nov. 2017, doi: 10.1177/0269215517700696.

[36]

L. R. Nascimento, S. M. Michaelsen, L. Ada, J. C. Polese, and L. F. Teixeira-Salmela, 'Cyclical electrical stimulation increases strength and improves activity after stroke: a systematic review', *Journal of Physiotherapy*, vol. 60, no. 1, pp. 22–30, Mar. 2014, doi: 10.1016/j.jphys.2013.12.002.

[37]

A. Pollock et al., 'Interventions for improving upper limb function after stroke', *Cochrane Database of Systematic Reviews*, Nov. 2014, doi: 10.1002/14651858.CD010820.pub2.

[38]

D. N. Rushton, 'Functional Electrical Stimulation and rehabilitation—an hypothesis', *Medical Engineering & Physics*, vol. 25, no. 1, pp. 75–78, Jan. 2003, doi: 10.1016/S1350-4533(02)00040-1.

[39]

C. Stein, C. G. Fritsch, C. Robinson, G. Sbruzzi, and R. D. M. Plentz, 'Effects of Electrical Stimulation in Spastic Muscles After Stroke', *Stroke*, vol. 46, no. 8, pp. 2197–2205, Aug. 2015, doi: 10.1161/STROKEAHA.115.009633.

[40]

A. K. Vafadar, J. N. Côté, and P. S. Archambault, 'Effectiveness of Functional Electrical Stimulation in Improving Clinical Outcomes in the Upper Arm following Stroke: A Systematic Review and Meta-Analysis', *BioMed Research International*, vol. 2015, pp. 1–14, 2015, doi: 10.1155/2015/729768.

[41]

J. M. Veerbeek et al., 'What Is the Evidence for Physical Therapy Poststroke? A Systematic Review and Meta-Analysis', *PLoS ONE*, vol. 9, no. 2, Feb. 2014, doi: 10.1371/journal.pone.0087987.

[42]

'The national service framework for long term conditions'. [Online]. Available: <https://www.gov.uk/government/publications/quality-standards-for-supporting-people-with-long-term-conditions>

[43]

K. James, *The strands of speech and language therapy: weaving a therapy plan for neurorehabilitation*. London: Speechmark, 2011.

[44]

M. Ylvisaker and T. Feeney, 'Reflections on Dobermanns, poodles, and social rehabilitation for difficult-to-serve individuals with traumatic brain injury', *Aphasiology*, vol. 14, no. 4, pp. 407–431, Apr. 2000, doi: 10.1080/026870300401432.

[45]

M. Ylvisaker and T. Feeney, 'Reconstruction of Identity After Brain Injury', *Brain Impairment*, vol. 1, no. 01, pp. 12–28, May 2000, doi: 10.1375/brim.1.1.12.

[46]

R. C. Katz, Brooke Hallowell, Chris Co, 'A multinational comparison of aphasia management practices', *International Journal of Language & Communication Disorders*, vol. 35, no. 2, pp. 303–314, Jan. 2000, doi: 10.1080/136828200247205.

[47]

A. L. Holland, *Language disorders in adults: recent advances*. San Diego, Calif: College-Hill Press, 1984.

[48]

J. A. Kleim and T. A. Jones, 'Principles of Experience-Dependent Neural Plasticity: Implications for Rehabilitation After Brain Damage', *Journal of Speech Language and Hearing Research*, vol. 51, no. 1, Feb. 2008, doi: 10.1044/1092-4388(2008/018).

[49]

'Journal of Rehabilitation Medicine - Abstract - The arm studio to intensify the upper limb rehabilitation after stroke: Concept, acceptance, utilization and preliminary clinical results'. [Online]. Available: <https://medicaljournals.se/jrm/content/abstract/10.2340/16501977-0517>

[50]

W. Liao, C. Wu, Y. Hsieh, K. Lin, and W. Chang, 'Effects of robot-assisted upper limb rehabilitation on daily function and real-world arm activity in patients with chronic stroke: a randomized controlled trial', *Clinical Rehabilitation*, vol. 26, no. 2, pp. 111–120, Feb. 2012, doi: 10.1177/0269215511416383.

[51]

'Journal of Rehabilitation Medicine - Abstract - Evaluation of functional outcome measures for the hemiparetic upper limb: A systematic review'. [Online]. Available: <https://www.medicaljournals.se/jrm/content/abstract/10.2340/16501977-0276>

[52]

K. Baker, S. J. Cano, and E. D. Playford, 'Outcome Measurement in Stroke: A Scale Selection Strategy', *Stroke*, vol. 42, no. 6, pp. 1787–1794, Jun. 2011, doi: 10.1161/STROKEAHA.110.608505.

[53]

R. Buchbinder, S. Green, J. M. Youd, R. V. Johnston, and M. Cumpston, 'Arthrographic distension for adhesive capsulitis (frozen shoulder)', *Cochrane Database of Systematic Reviews*, Jan. 2008, doi: 10.1002/14651858.CD007005.

[54]

Subhasish Chatterjee, 'The California Tri-pull Taping Method in the Treatment of Shoulder Subluxation After Stroke: A Randomized Clinical Trial', *North American Journal of Medical Sciences*, vol. 8, no. 4, 2016, doi: 10.4103/1947-2714.179933. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4866473/>

[55]

J. Chae et al., 'Poststroke Shoulder Pain: Its Relationship to Motor Impairment, Activity Limitation, and Quality of Life', *Archives of Physical Medicine and Rehabilitation*, vol. 88, no. 3, pp. 298–301, Mar. 2007, doi: 10.1016/j.apmr.2006.12.007.

[56]

G. E. Gamble, E. Barberan, D. Bowsher, P. J. Tyrrell, and A. K. P. Jones, 'Post stroke shoulder pain: more common than previously realized', *European Journal of Pain*, vol. 4, no. 3, pp. 313–315, Sep. 2000, doi: 10.1053/eujp.2000.0192.

[57]

S. Green, R. Buchbinder, and S. E. Hetrick, 'Physiotherapy interventions for shoulder pain', *Cochrane Database of Systematic Reviews*, Apr. 2003, doi: 10.1002/14651858.CD004258.

[58]

'Effects of neuromuscular electrical stimulation on arterial hemodynamic properties and body composition in paretic upper extremities of patients with subacute stroke'. [Online]. Available: http://biomedj.cgu.edu.tw/pdfs/2014/37/4/images/BiomedJ_2014_37_4_205_117892.pdf

[59]

'Use of an integrated care pathway: a third round audit of the management of shoulder pain in neurological conditions'. [Online]. Available: <https://medicaljournals.se/jrm/content/abstract/10.1080/16501970310012446>

[60]

S. McDonald, L. Togher, and C. Code, Social and communication disorders following traumatic brain injury, Second edition. London: Psychology Press, Taylor & Francis Group, 2014.

[61]

C. A. Mateer, C. S. Sira, and M. E. O'Connell, 'Putting Humpty Dumpty Together Again', *Journal of Head Trauma Rehabilitation*, vol. 20, no. 1, pp. 62–75, Jan. 2005, doi: 10.1097/00001199-200501000-00007.

[62]

S. Tornås et al., 'Rehabilitation of Executive Functions in Patients with Chronic Acquired Brain Injury with Goal Management Training, External Cuing, and Emotional Regulation: A Randomized Controlled Trial', *Journal of the International Neuropsychological Society*, vol. 22, no. 04, pp. 436–452, Apr. 2016, doi: 10.1017/S1355617715001344.

[63]

T. M. McMillan and R. L. Wood, Eds., *Neurobehavioural disability and social handicap following traumatic brain injury*, Second edition. London: Routledge, 2017.

[64]

J. Winegardner, C. Keohane, L. Prince, and D. Neumann, 'Perspective training to treat anger problems after brain injury: Two case studies', *NeuroRehabilitation*, vol. 39, no. 1, pp. 153–162, Jul. 2016, doi: 10.3233/NRE-161347.

[65]

L. Cipolotti and E. K. Warrington, 'Neuropsychological assessment.', *Journal of Neurology, Neurosurgery & Psychiatry*, vol. 58, no. 6, pp. 655–664, Jun. 1995, doi: 10.1136/jnnp.58.6.655.

[66]

P. S. Sachdev, D. M. Lipnicki, J. D. Crawford, W. Wen, and H. Brodaty, 'Progression of cognitive impairment in stroke/TIA patients over 3 years', *Journal of Neurology, Neurosurgery & Psychiatry*, vol. 85, no. 12, pp. 1324–1330, Dec. 2014, doi: 10.1136/jnnp-2013-306776.

[67]

G. M. S. Nys et al., 'The prognostic value of domain-specific cognitive abilities in acute first-ever stroke', *Neurology*, vol. 64, no. 5, pp. 821–827, Mar. 2005, doi: 10.1212/01.WNL.0000152984.28420.5A.

[68]

R. Hurford, A. Charidimou, Z. Fox, L. Cipolotti, and D. J. Werring, 'Domain-specific trends in cognitive impairment after acute ischaemic stroke', *Journal of Neurology*, vol. 260, no. 1, pp. 237–241, Jan. 2013, doi: 10.1007/s00415-012-6625-0.

[69]

C. M. Van Heugten, L. Walton, and U. Hentschel, 'Can we forget the Mini-Mental State Examination? A systematic review of the validity of cognitive screening instruments within one month after stroke', *Clinical Rehabilitation*, vol. 29, no. 7, pp. 694–704, Jul. 2015, doi: 10.1177/0269215514553012.

[70]

'Manually add a new bookmark | University College London'. [Online]. Available: <http://readinglists.ucl.ac.uk/ui/forms/bookmarklet.html?fast=true&title=Journal%20of%20Rehabilitation%20Medicine%20-%20Abstract%20-%20Screening%20for%20cognitive%20impairment%20after%20stroke%3A%20A%20systematic%20review%20of%20psychometric%20properties%20and%20clinical%20utility&uri=https%253A%252F%252Fwww.medicaljournals.se%252Fjrm%252Fcontent%252Fabstract%252F10.2340%252F165019>

77-1930

[71]

E. Chan, S. Altendorff, C. Healy, D. J. Werring, and L. Cipolotti, 'The test accuracy of the Montreal Cognitive Assessment (MoCA) by stroke lateralisation', *Journal of the Neurological Sciences*, vol. 373, pp. 100–104, Feb. 2017, doi: 10.1016/j.jns.2016.12.028.

[72]

E. Chan, S. Khan, R. Oliver, S. K. Gill, D. J. Werring, and L. Cipolotti, 'Underestimation of cognitive impairments by the Montreal Cognitive Assessment (MoCA) in an acute stroke unit population', *Journal of the Neurological Sciences*, vol. 343, no. 1–2, pp. 176–179, Aug. 2014, doi: 10.1016/j.jns.2014.05.005.

[73]

P. Craig, P. Dieppe, S. Macintyre, S. Michie, I. Nazareth, and M. Petticrew, 'Developing and evaluating complex interventions: the new Medical Research Council guidance', *BMJ*, Sep. 2008, doi: 10.1136/bmj.a1655.

[74]

R. C. Katz and R. T. Wertz, 'The Efficacy of Computer-Provided Reading Treatment for Chronic Aphasic Adults', *Journal of Speech Language and Hearing Research*, vol. 40, no. 3, Jun. 1997, doi: 10.1044/jslhr.4003.493.

[75]

J. Marshall et al., 'Evaluating the Benefits of Aphasia Intervention Delivered in Virtual Reality: Results of a Quasi-Randomised Study', *PLOS ONE*, vol. 11, no. 8, Aug. 2016, doi: 10.1371/journal.pone.0160381.

[76]

R. Palmer et al., 'Computer Therapy Compared With Usual Care for People With Long-Standing Aphasia Poststroke: A Pilot Randomized Controlled Trial', *Stroke*, vol. 43, no. 7, pp. 1904–1911, Jul. 2012, doi: 10.1161/STROKEAHA.112.650671.

[77]

R. Varley, P. E. Cowell, L. Dyson, L. Inglis, A. Roper, and S. P. Whiteside, 'Self-Administered Computer Therapy for Apraxia of Speech', *Stroke*, Jan. 2016, doi: 10.1161/STROKEAHA.115.011939.

[78]

A. Bowen, C. Hazelton, A. Pollock, and N. B. Lincoln, 'Cognitive rehabilitation for spatial neglect following stroke', *Cochrane Database of Systematic Reviews*, Jul. 2013, doi: 10.1002/14651858.CD003586.pub3.

[79]

M. Corbetta, M. J. Kincade, C. Lewis, A. Z. Snyder, and A. Sapir, 'Neural basis and recovery of spatial attention deficits in spatial neglect', *Nature Neuroscience*, vol. 8, no. 11, pp. 1603–1610, Nov. 2005, doi: 10.1038/nn1574.

[80]

J. M. Ferro, G. Mariano, and S. Madureira, 'Recovery from Aphasia and Neglect', *Cerebrovascular Diseases*, vol. 9, no. Suppl. 5, pp. 6–22, 1999, doi: 10.1159/000047571.

[81]

F. Frassinetti, V. Angeli, F. Meneghello, S. Avanzi, and E. Ladavas, 'Long-lasting amelioration of visuospatial neglect by prism adaptation', *Brain*, vol. 125, no. 3, pp. 608–623, Mar. 2002, doi: 10.1093/brain/awf056.

[82]

N. Gorgoraptis et al., 'The effects of the dopamine agonist rotigotine on hemispatial neglect following stroke', *Brain*, vol. 135, no. 8, pp. 2478–2491, Aug. 2012, doi: 10.1093/brain/aws154.

[83]

R. F. Kaplan, M. Verfaellie, M.-E. Meadows, L. R. Caplan, M. S. Pessin, and L. D. DeWitt, 'Changing Attentional Demands in Left Hemispatial Neglect', *Archives of Neurology*, vol.

48, no. 12, pp. 1263–1266, Dec. 1991, doi: 10.1001/archneur.1991.00530240067023.

[84]

G. Kerkhoff et al., 'Smooth Pursuit "Bedside" Training Reduces Disability and Unawareness During the Activities of Daily Living in Neglect', *Neurorehabilitation and Neural Repair*, vol. 28, no. 6, pp. 554–563, Jul. 2014, doi: 10.1177/1545968313517757.

[85]

N. Koiava, Y.-H. Ong, M. M. Brown, J. Acheson, G. T. Plant, and A. P. Leff, 'A "web app" for diagnosing hemianopia', *Journal of Neurology, Neurosurgery & Psychiatry*, vol. 83, no. 12, pp. 1222–1224, Dec. 2012, doi: 10.1136/jnnp-2012-302270.

[86]

J. I. Grieve, *Neuropsychology for occupational therapists: cognition in occupational performance*, Fourth edition. Hoboken, NJ, USA: Wiley Blackwell, 2017.

[87]

G. Gillen and St Bartholomew School of Nursing and Midwifery, *Cognitive and perceptual rehabilitation: optimizing function*. St. Louis, Mo: Mosby/Elsevier, 2009.

[88]

Amy J. Bastian, 'Understanding sensorimotor adaptation and learning for rehabilitation', *Current opinion in neurology*, vol. 21, no. 6, 2008, doi: 10.1097/WCO.0b013e328315a293. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2954436/>

[89]

J. Bernhardt, M. N. Thuy, J. M. Collier, and L. A. Legg, 'Very early versus delayed mobilisation after stroke', *Cochrane Database of Systematic Reviews*, Jan. 2009, doi: 10.1002/14651858.CD006187.pub2.

[90]

'Muscle strength and muscle training after stroke'. [Online]. Available: <https://www.medicaljournals.se/jrm/content/abstract/10.2340/16501977-0018>

[91]

L. Carey, R. Macdonell, and T. A. Matyas, 'SENSe: Study of the Effectiveness of Neurorehabilitation on Sensation', *Neurorehabilitation and Neural Repair*, vol. 25, no. 4, pp. 304–313, May 2011, doi: 10.1177/1545968310397705.

[92]

L. Connell and S. Tyson, 'Measures of sensation in neurological conditions: a systematic review', *Clinical Rehabilitation*, vol. 26, no. 1, pp. 68–80, Jan. 2012, doi: 10.1177/0269215511412982.

[93]

L. A. Connell, N. E. McMahon, and N. Adams, 'Stroke survivors' experiences of somatosensory impairment after stroke: An Interpretative Phenomenological Analysis', *Physiotherapy*, vol. 100, no. 2, pp. 150–155, Jun. 2014, doi: 10.1016/j.physio.2013.09.003.

[94]

M. Demetrios, F. Khan, L. Turner-Stokes, C. Brand, and S. McSweeney, 'Multidisciplinary rehabilitation following botulinum toxin and other focal intramuscular treatment for post-stroke spasticity', *Cochrane Database of Systematic Reviews*, Jun. 2013, doi: 10.1002/14651858.CD009689.pub2.

[95]

S. Doyle, S. Bennett, S. E. Fasoli, and K. T. McKenna, 'Interventions for sensory impairment in the upper limb after stroke', *Cochrane Database of Systematic Reviews*, Jun. 2010, doi: 10.1002/14651858.CD006331.pub2.

[96]

Klemens Fheodoroff, 'Factors Influencing Goal Attainment in Patients with Post-Stroke Upper Limb Spasticity Following Treatment with Botulinum Toxin A in Real-Life Clinical Practice: Sub-Analyses from the Upper Limb International Spasticity (ULIS)-II Study', *Toxins*

, vol. 7, no. 4, 2015, doi: 10.3390/toxins7041192. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4417963/>