

GEOGG065: Environmental GIS

View Online



[1]

Baily, B. and Nowell, D. 1996. Techniques for monitoring coastal change: a review and case study. *Ocean & Coastal Management*. 32, 2 (1996), 85–95.
[https://doi.org/10.1016/S0964-5691\(96\)00058-0](https://doi.org/10.1016/S0964-5691(96)00058-0).

[2]

Band, L.E. Topographic Partition of Watersheds with Digital Elevation Models. *Water Resources Research*. 22, 1, 15–24. <https://doi.org/10.1029/WR022i001p00015>.

[3]

Barfoot, P.J. and Tucker, J.J. 1980. Geomorphological changes at Blakeney Point. *Transactions of the Norfolk and Norwich Naturalists Society*. 25, 2 (1980), 49–60.

[4]

Bernhardsen, T. 1999. *Geographic information systems: an introduction*. Wiley.

[5]

Bock, M. 2003. Remote sensing and GIS-based techniques for the classification and monitoring of biotopes: Case examples for a wet grass- and moor land area in Northern Germany. *Journal for Nature Conservation*. 11, 3 (2003), 145–155.
<https://doi.org/10.1078/1617-1381-00050>.

[6]

Boteva, D. et al. Evaluation and mapping of the conservation significance of habitats using GIS: an example from Crete, Greece. *Journal for Nature Conservation*. 12, 4, 237–250.
<https://doi.org/10.1016/j.jnc.2004.09.002>.

[7]

Brewer, C.A. 2005. *Designing better maps: a guide for GIS users*. ESRI Press.

[8]

Carmona, C.P. et al. Grazing management or physiography? Factors controlling vegetation recovery in Mediterranean grasslands. *Ecological Modelling*. 251, 73–84.
<https://doi.org/10.1016/j.ecolmodel.2012.12.005>.

[9]

Chaaban, F. and et al 2012. Using ArcGISH Modelbuilder and Aerial Photographs to Measure Coastline Retreat and Advance: North of France. *Journal of Coastal Research*. 28, 6 (2012), 1567–1579.

[10]

Chen, C.-Y. and Yu, F.-C. Morphometric analysis of debris flows and their source areas using GIS. *Geomorphology*. 129, 3–4, 387–397.
<https://doi.org/10.1016/j.geomorph.2011.03.002>.

[11]

Dalton, T. et al. Mapping human dimensions in marine spatial planning and management: An example from Narragansett Bay, Rhode Island. *Marine Policy*. 34, 2, 309–319.
<https://doi.org/10.1016/j.marpol.2009.08.001>.

[12]

DeMers, M.N. 2009. *Fundamentals of geographic information systems*. Wiley.

[13]

Dong, Z. et al. Monitoring sand dune advance in the Taklimakan Desert. *Geomorphology*. 35, 3-4, 219-231. [https://doi.org/10.1016/S0169-555X\(00\)00039-8](https://doi.org/10.1016/S0169-555X(00)00039-8).

[14]

Draper, D. and et al Application of GIS in plant conservation programmes in Portugal. *Biological Conservation*. 113, 3, 337-349. [https://doi.org/10.1016/S0006-3207\(03\)00125-3](https://doi.org/10.1016/S0006-3207(03)00125-3).

[15]

Duane Nellis, M. et al. Remote sensing of temporal and spatial variations in pool size, suspended sediment, turbidity, and Secchi depth in Tuttle Creek Reservoir, Kansas: 1993. *Geomorphology*. 21, 3-4, 281-293. [https://doi.org/10.1016/S0169-555X\(97\)00067-6](https://doi.org/10.1016/S0169-555X(97)00067-6).

[16]

Ekeboom, J. and Erkkila, A. Using aerial photography for identification of marine and coastal habitats under the EU's Habitats Directive. *Aquatic Conservation: Marine and Freshwater Ecosystems*. 13, 4, 287-304. <https://doi.org/10.1002/aqc.553>.

[17]

Estes, J.E. 1966. Some applications of aerial infrared imagery. Allen Press Inc.

[18]

Evaluation of Riparian Buffer Zones using GIS and Remote Sensing to Target Watershed Restoration Efforts: 2004. <http://igs.indiana.edu/WatershedHydrology/RiparianBuffers.cfm>.

[19]

Fromard, F. et al. Half a century of dynamic coastal change affecting mangrove shorelines of French Guiana. A case study based on remote sensing data analyses and field surveys. *Marine Geology*. 208, 2-4, 265-280. <https://doi.org/10.1016/j.margeo.2004.04.018>.

[20]

Geospatial Analysis: 2021. <https://www.spatialanalysisonline.com/HTML/index.html>.

[21]

Greco, S.E. and Plant, R.E. Temporal mapping of riparian landscape change on the Sacramento river, miles 196–218, California, USA. *Landscape Research*. 28, 4, 405–426. <https://doi.org/10.1080/0142639032000150149>.

[22]

Hall, S.T. and Post, C.J. 2008. Advanced GIS Exercise: Estimating Beach and Dune Erosion in Coastal South Carolina. *Journal of Natural Resources and Life Sciences Education*. 37, (2008), 49–52.

[23]

Heywood, I. et al. 2006. *An introduction to geographical information systems*. Pearson Education.

[24]

Kadmon, R. and Harari-Kremer, R. Studying Long-Term Vegetation Dynamics Using Digital Processing of Historical Aerial Photographs. *Remote Sensing of Environment*. 68, 2, 164–176. [https://doi.org/10.1016/S0034-4257\(98\)00109-6](https://doi.org/10.1016/S0034-4257(98)00109-6).

[25]

Kienzle, S. The Effect of DEM Raster Resolution on First Order, Second Order and Compound Terrain Derivatives. *Transactions in GIS*. 8, 1, 83–111. <https://doi.org/10.1111/j.1467-9671.2004.00169.x>.

[26]

Koch, T. 2004. *The Map as Intent: Variations on the Theme of John Snow*. *Cartographica*.

[27]

Leeks, G.J.L. and et al The LOIS river monitoring network: strategy and implementation. *Science of The Total Environment*. 194-195, 101-109.
[https://doi.org/10.1016/S0048-9697\(96\)05356-9](https://doi.org/10.1016/S0048-9697(96)05356-9).

[28]

Leonard, P.B. and et al Remote detection of small wetlands in the Atlantic coastal plain of North America: Local relief models, ground validation, and high-throughput computing. *Forest Ecology and Management*. 284, 107-115.
<https://doi.org/10.1016/j.foreco.2012.07.034>.

[29]

Longley, P. and et al 2011. *Geographic information systems & science*. John Wiley & Sons.

[30]

Lundblad, E.R. and et al 2006. A Benthic Terrain Classification Scheme for American Samoa. *Marine Geodesy*. 29, 2 (2006), 89-111.
<https://doi.org/10.1080/01490410600738021>.

[31]

Maddrell, R.J. Managed coastal retreat, reducing flood risks and protection costs, Dungeness Nuclear Power Station, UK. *Coastal Engineering*. 28, 1-4, 1-15.
[https://doi.org/10.1016/0378-3839\(95\)00035-6](https://doi.org/10.1016/0378-3839(95)00035-6).

[32]

Manson, F.J. et al. Spatial and temporal variation in distribution of mangroves in Moreton Bay, subtropical Australia: a comparison of pattern metrics and change detection analyses based on aerial photographs. *Estuarine, Coastal and Shelf Science*. 57, 4, 653-666.
[https://doi.org/10.1016/S0272-7714\(02\)00405-5](https://doi.org/10.1016/S0272-7714(02)00405-5).

[33]

McGregor, S.J. An integrated geographic information system approach for modeling the suitability of conifer habitat in an alpine environment. *Geomorphology*. 21, 3-4, 265-280.
[https://doi.org/10.1016/S0169-555X\(97\)00066-4](https://doi.org/10.1016/S0169-555X(97)00066-4).

[34]

McKenna et al., J. 2003. Obsolete Maps and Coastal Management: Case Studies from Northwest Ireland. *Coastal Management*. 31, 3 (2003), 229–246.

[35]

Meso-scale morphological changes in the Loughros More estuary: 2002.
<http://sheskinmore.wordpress.com/research/>.

[36]

van der Meulen, F. et al. The use of a GIS in assessing the impacts of sea level rise on nature conservation along the Dutch coast: 1990–2090. *Landscape Ecology*. 6, 1–2, 105–113. <https://doi.org/10.1007/BF00157750>.

[37]

Miller, S.N. and et al The Automated Geospatial Watershed Assessment tool. *Environmental Modelling & Software*. 22, 3, 365–377.
<https://doi.org/10.1016/j.envsoft.2005.12.004>.

[38]

Moore, T. and et al An Expert System for Integrated Coastal Zone Management: A Geomorphological Case Study. *Marine Pollution Bulletin*. 37, 3–7, 361–370.
[https://doi.org/10.1016/S0025-326X\(99\)00129-0](https://doi.org/10.1016/S0025-326X(99)00129-0).

[39]

Multibeam sonar mapping and modeling of a submerged bryophyte mat in Crater Lake, Oregon: 2008. <http://pubs.usgs.gov/ds/366/>.

[40]

Naito, A.T. and Cairns, D.M. Relationships between Arctic shrub dynamics and topographically derived hydrologic characteristics. *Environmental Research Letters*. 6, 4.
<https://doi.org/10.1088/1748-9326/6/4/045506>.

[41]

Nams, V.O. et al. Determining the spatial scale for conservation purposes – an example with grizzly bears. *Biological Conservation*. 128, 1, 109–119.
<https://doi.org/10.1016/j.biocon.2005.09.020>.

[42]

Neilson, B. and Costello, M.J. The Relative Lengths of Seashore Substrata Around the Coastline of Ireland as Determined by Digital Methods in a Geographical Information System. *Estuarine, Coastal and Shelf Science*. 49, 4, 501–508.
<https://doi.org/10.1006/ecss.1999.0507>.

[43]

Pedersen, Å.Ø. et al. Implementation of a GIS-based management tool for conservation of biodiversity within the municipality of Oslo, Norway. *Landscape and Urban Planning*. 68, 4, 429–438. [https://doi.org/10.1016/S0169-2046\(03\)00148-8](https://doi.org/10.1016/S0169-2046(03)00148-8).

[44]

Perotto-Baldiviezo, H.L. and et al GIS-based spatial analysis and modeling for landslide hazard assessment in steeplands, southern Honduras. *Agriculture, Ecosystems & Environment*. 103, 1, 165–176. <https://doi.org/10.1016/j.agee.2003.10.011>.

[45]

Rozo, M.G. et al. The anastomosing pattern and the extensively distributed scroll bars in the middle Amazon River. *Earth Surface Processes and Landforms*. 37, 14, 1471–1488.
<https://doi.org/10.1002/esp.3249>.

[46]

Rumsby, B. Vertical accretion rates in fluvial systems: -a comparison of volumetric and depth-based estimates. *Earth Surface Processes and Landforms*. 25, 6, 617–631.
[https://doi.org/10.1002/1096-9837\(200006\)25:6<617::AID-ESP99>3.0.CO;2-Z](https://doi.org/10.1002/1096-9837(200006)25:6<617::AID-ESP99>3.0.CO;2-Z).

[47]

Rumsby, B.T. and Macklin, M.G. Channel and floodplain response to recent abrupt climate change: The tyne basin, Northern England. *Earth Surface Processes and Landforms*. 19, 6, 499–515. <https://doi.org/10.1002/esp.3290190603>.

[48]

Salem, B.B. Application of GIS to biodiversity monitoring. *Journal of Arid Environments*. 54, 1, 91–114. <https://doi.org/10.1006/jare.2001.0887>.

[49]

Sear, D.A. and Newson, M.D. Environmental change in river channels: a neglected element. Towards geomorphological typologies, standards and monitoring. *Science of The Total Environment*. 310, 1–3, 17–23. [https://doi.org/10.1016/S0048-9697\(02\)00619-8](https://doi.org/10.1016/S0048-9697(02)00619-8).

[50]

Sheppard, C.R.C. and et al Habitat mapping in the Caribbean for management and conservation: Use and assessment of aerial photography. *Aquatic Conservation: Marine and Freshwater Ecosystems*. 5, 4, 277–298. <https://doi.org/10.1002/aqc.3270050404>.

[51]

Sørensen, R. et al. On the calculation of the topographic wetness index: evaluation of different methods based on field observations. *Hydrology and Earth System Sciences*. 10, 1, 101–112. <https://doi.org/10.5194/hess-10-101-2006>.

[52]

Sutherland, J. Error analysis of Ordnance Survey map tidelines, UK. *Proceedings of the ICE - Maritime Engineering*. 165, 4, 189–197. <https://doi.org/10.1680/maen.2011.10>.

[53]

Tarboton, D.G. et al. On the extraction of channel networks from digital elevation data. *Hydrological Processes*. 5, 1, 81–100. <https://doi.org/10.1002/hyp.3360050107>.

[54]

Taylor, J.C. et al. Monitoring landscape change in the National Parks of England and Wales using aerial photo interpretation and GIS. *International Journal of Remote Sensing*. 21, 13–14, 2737–2752. <https://doi.org/10.1080/01431160050110269>.

[55]

Teng, J. et al. A GIS-Based Tool for Spatial and Distributed Hydrological Modelling: CLASS Spatial Analyst. *Transactions in GIS*. 12, 2, 209–225. <https://doi.org/10.1111/j.1467-9671.2008.01096.x>.

[56]

Thompson, S. and et al Identifying potential breeding sites for the stone curlew (*Burhinus oedicephalus*) in the UK. *Journal for Nature Conservation*. 12, 4, 229–235. <https://doi.org/10.1016/j.jnc.2004.07.002>.

[57]

Thorbjarnarson, J. and et al Regional habitat conservation priorities for the American crocodile. *Biological Conservation*. 128, 1, 25–36. <https://doi.org/10.1016/j.biocon.2005.09.013>.

[58]

Vitek, J.D. et al. Mapping geomorphology: A journey from paper maps, through computer mapping to GIS and Virtual Reality. *Geomorphology*. 16, 3, 233–249. [https://doi.org/10.1016/S0169-555X\(96\)80003-1](https://doi.org/10.1016/S0169-555X(96)80003-1).

[59]

Walsh, S.J. et al. An overview of scale, pattern, process relationships in geomorphology: a remote sensing and GIS perspective. *Geomorphology*. 21, 3–4, 183–205. [https://doi.org/10.1016/S0169-555X\(97\)00057-3](https://doi.org/10.1016/S0169-555X(97)00057-3).

[60]

White, K. and El Asmar, H.M. Monitoring changing position of coastlines using Thematic Mapper imagery, an example from the Nile Delta. *Geomorphology*. 29, 1–2, 93–105. [https://doi.org/10.1016/S0169-555X\(99\)00008-2](https://doi.org/10.1016/S0169-555X(99)00008-2).

[61]

Wilson, M.F.J. and et al Multiscale Terrain Analysis of Multibeam Bathymetry Data for Habitat Mapping on the Continental Slope. *Marine Geodesy*. 30, 1-2, 3-35.
<https://doi.org/10.1080/01490410701295962>.

[62]

Wright, D.J. and Heyman, W.D. Introduction to the Special Issue: Marine and Coastal GIS for Geomorphology, Habitat Mapping, and Marine Reserves. *Marine Geodesy*. 31, 4, 223-230. <https://doi.org/10.1080/01490410802466306>.