

GEOGG065: Environmental GIS

View Online



[1]

C. A. Brewer, *Designing better maps: a guide for GIS users*. Redlands, Calif: ESRI Press, 2005.

[2]

M. N. DeMers, *Fundamentals of geographic information systems*, 4th ed. Hoboken, NJ: Wiley, 2009.

[3]

M. de Smith, P. Longley, and M. Goodchild, 'Geospatial Analysis', 2021. Available: <https://www.spatialanalysisonline.com/HTML/index.html>

[4]

I. Heywood, S. Cornelius, and S. Carver, *An introduction to geographical information systems*, 3rd ed. Harlow: Pearson Education, 2006.

[5]

P. Longley and et al, *Geographic information systems & science*, 3rd ed. Hoboken, N.J.: John Wiley & Sons, 2011.

[6]

B. Baily and D. Nowell, 'Techniques for monitoring coastal change: a review and case study', *Ocean & Coastal Management*, vol. 32, no. 2, pp. 85–95, 1996, doi:

10.1016/S0964-5691(96)00058-0

[7]

P. J. Barfoot and J. J. Tucker, 'Geomorphological changes at Blakeney Point', Transactions of the Norfolk and Norwich Naturalists Society, vol. 25, no. 2, pp. 49-60, 1980.

[8]

T. Bernhardsen, Geographic information systems: an introduction, 2nd ed. New York: Wiley, 1999.

[9]

M. Bock, 'Remote sensing and GIS-based techniques for the classification and monitoring of biotopes: Case examples for a wet grass- and moor land area in Northern Germany', Journal for Nature Conservation, vol. 11, no. 3, pp. 145-155, 2003, doi: 10.1078/1617-1381-00050

[10]

D. Boteva, G. Griffiths, and P. Dimopoulos, 'Evaluation and mapping of the conservation significance of habitats using GIS: an example from Crete, Greece', Journal for Nature Conservation, vol. 12, no. 4, pp. 237-250, doi: 10.1016/j.jnc.2004.09.002

[11]

H. Burningham, 'Meso-scale morphological changes in the Loughros More estuary', Proceedings of Littoral 2002: The Changing Coast, 2002. Available: <http://sheskinmore.wordpress.com/research/>

[12]

Z. Dong, X. Wang, and G. Chen, 'Monitoring sand dune advance in the Taklimakan Desert', Geomorphology, vol. 35, no. 3-4, pp. 219-231, doi: 10.1016/S0169-555X(00)00039-8

[13]

D. Draper and et al, 'Application of GIS in plant conservation programmes in Portugal', *Biological Conservation*, vol. 113, no. 3, pp. 337–349, doi: 10.1016/S0006-3207(03)00125-3

[14]

J. Ekebom and A. Erkkila, 'Using aerial photography for identification of marine and coastal habitats under the EU's Habitats Directive', *Aquatic Conservation: Marine and Freshwater Ecosystems*, vol. 13, no. 4, pp. 287–304, doi: 10.1002/aqc.553

[15]

F. Fromard, C. Vega, and C. Proisy, 'Half a century of dynamic coastal change affecting mangrove shorelines of French Guiana. A case study based on remote sensing data analyses and field surveys', *Marine Geology*, vol. 208, no. 2–4, pp. 265–280, doi: 10.1016/j.margeo.2004.04.018

[16]

G. J. L. Leeks and et al, 'The LOIS river monitoring network: strategy and implementation', *Science of The Total Environment*, vol. 194–195, pp. 101–109, doi: 10.1016/S0048-9697(96)05356-9

[17]

S. L. Letsinger, 'Evaluation of Riparian Buffer Zones using GIS and Remote Sensing to Target Watershed Restoration Efforts', 2004. Available: <http://igs.indiana.edu/WatershedHydrology/RiparianBuffers.cfm>

[18]

T. Koch, 'The Map as Intent: Variations on the Theme of John Snow', *Cartographica*, vol. 39, no. 4, pp. 1–15, 2004. Available: [http://www.ph.ucla.edu/epi/snow/cartographica39\(4\)1_14_2004.pdf](http://www.ph.ucla.edu/epi/snow/cartographica39(4)1_14_2004.pdf)

[19]

R. J. Maddrell, 'Managed coastal retreat, reducing flood risks and protection costs, Dungeness Nuclear Power Station, UK', *Coastal Engineering*, vol. 28, no. 1–4, pp. 1–15, doi: 10.1016/0378-3839(95)00035-6

[20]

F. J. Manson, N. R. Loneragan, and S. R. Phinn, 'Spatial and temporal variation in distribution of mangroves in Moreton Bay, subtropical Australia: a comparison of pattern metrics and change detection analyses based on aerial photographs', *Estuarine, Coastal and Shelf Science*, vol. 57, no. 4, pp. 653–666, doi: 10.1016/S0272-7714(02)00405-5

[21]

S. J. McGregor, 'An integrated geographic information system approach for modeling the suitability of conifer habitat in an alpine environment', *Geomorphology*, vol. 21, no. 3–4, pp. 265–280, doi: 10.1016/S0169-555X(97)00066-4

[22]

J. McKenna et al., 'Obsolete Maps and Coastal Management: Case Studies from Northwest Ireland', *Coastal Management*, vol. 31, no. 3, pp. 229–246, 2003, Available: <http://search.ebscohost.com/login.aspx?direct=true&AuthType=ip,shib&db=bth&AN=9930623&site=ehost-live&scope=site>

[23]

T. Moore and et al, 'An Expert System for Integrated Coastal Zone Management: A Geomorphological Case Study', *Marine Pollution Bulletin*, vol. 37, no. 3–7, pp. 361–370, doi: 10.1016/S0025-326X(99)00129-0

[24]

V. O. Nams, G. Mowat, and M. A. Panian, 'Determining the spatial scale for conservation purposes – an example with grizzly bears', *Biological Conservation*, vol. 128, no. 1, pp. 109–119, doi: 10.1016/j.biocon.2005.09.020

[25]

B. Neilson and M. J. Costello, 'The Relative Lengths of Seashore Substrata Around the Coastline of Ireland as Determined by Digital Methods in a Geographical Information System', *Estuarine, Coastal and Shelf Science*, vol. 49, no. 4, pp. 501–508, doi: 10.1006/ecss.1999.0507

[26]

M. Duane Nellis, J. A. Harrington, and J. Wu, 'Remote sensing of temporal and spatial variations in pool size, suspended sediment, turbidity, and Secchi depth in Tuttle Creek Reservoir, Kansas: 1993', *Geomorphology*, vol. 21, no. 3-4, pp. 281-293, doi: 10.1016/S0169-555X(97)00067-6

[27]

Å. Ø. Pedersen, S. Nyhuus, T. Blindheim, and O. M. W. Krog, 'Implementation of a GIS-based management tool for conservation of biodiversity within the municipality of Oslo, Norway', *Landscape and Urban Planning*, vol. 68, no. 4, pp. 429-438, doi: 10.1016/S0169-2046(03)00148-8

[28]

H. L. Perotto-Baldiviezo and et al, 'GIS-based spatial analysis and modeling for landslide hazard assessment in steeplands, southern Honduras', *Agriculture, Ecosystems & Environment*, vol. 103, no. 1, pp. 165-176, doi: 10.1016/j.agee.2003.10.011

[29]

B. B. Salem, 'Application of GIS to biodiversity monitoring', *Journal of Arid Environments*, vol. 54, no. 1, pp. 91-114, doi: 10.1006/jare.2001.0887

[30]

D. A. Sear and M. D. Newson, 'Environmental change in river channels: a neglected element. Towards geomorphological typologies, standards and monitoring', *Science of The Total Environment*, vol. 310, no. 1-3, pp. 17-23, doi: 10.1016/S0048-9697(02)00619-8

[31]

S. Thompson and et al, 'Identifying potential breeding sites for the stone curlew (*Burhinus oedicnemus*) in the UK', *Journal for Nature Conservation*, vol. 12, no. 4, pp. 229-235, doi: 10.1016/j.jnc.2004.07.002

[32]

J. Thorbjarnarson and et al, 'Regional habitat conservation priorities for the American

crocodile', *Biological Conservation*, vol. 128, no. 1, pp. 25–36, doi: 10.1016/j.biocon.2005.09.013

[33]

S. J. Walsh, D. R. Butler, and G. P. Malanson, 'An overview of scale, pattern, process relationships in geomorphology: a remote sensing and GIS perspective', *Geomorphology*, vol. 21, no. 3–4, pp. 183–205, doi: 10.1016/S0169-555X(97)00057-3

[34]

K. White and H. M. El Asmar, 'Monitoring changing position of coastlines using Thematic Mapper imagery, an example from the Nile Delta', *Geomorphology*, vol. 29, no. 1–2, pp. 93–105, doi: 10.1016/S0169-555X(99)00008-2

[35]

F. van der Meulen, J. V. Witter, and S. M. Arens, 'The use of a GIS in assessing the impacts of sea level rise on nature conservation along the Dutch coast: 1990–2090', *Landscape Ecology*, vol. 6, no. 1–2, pp. 105–113, doi: 10.1007/BF00157750

[36]

J. D. Vitek, J. R. Giardino, and J. W. Fitzgerald, 'Mapping geomorphology: A journey from paper maps, through computer mapping to GIS and Virtual Reality', *Geomorphology*, vol. 16, no. 3, pp. 233–249, doi: 10.1016/S0169-555X(96)80003-1

[37]

L. E. Band, 'Topographic Partition of Watersheds with Digital Elevation Models', *Water Resources Research*, vol. 22, no. 1, pp. 15–24, doi: 10.1029/WR022i001p00015

[38]

C. P. Carmona, A. Röder, F. M. Azcárate, and B. Peco, 'Grazing management or physiography? Factors controlling vegetation recovery in Mediterranean grasslands', *Ecological Modelling*, vol. 251, pp. 73–84, doi: 10.1016/j.ecolmodel.2012.12.005

[39]

C.-Y. Chen and F.-C. Yu, 'Morphometric analysis of debris flows and their source areas using GIS', *Geomorphology*, vol. 129, no. 3–4, pp. 387–397, doi: 10.1016/j.geomorph.2011.03.002

[40]

T. Dalton, R. Thompson, and D. Jin, 'Mapping human dimensions in marine spatial planning and management: An example from Narragansett Bay, Rhode Island', *Marine Policy*, vol. 34, no. 2, pp. 309–319, doi: 10.1016/j.marpol.2009.08.001

[41]

P. Dartnell and et al, 'Multibeam sonar mapping and modeling of a submerged bryophyte mat in Crater Lake, Oregon', 2008. Available: <http://pubs.usgs.gov/ds/366/>

[42]

S. Kienzle, 'The Effect of DEM Raster Resolution on First Order, Second Order and Compound Terrain Derivatives', *Transactions in GIS*, vol. 8, no. 1, pp. 83–111, doi: 10.1111/j.1467-9671.2004.00169.x

[43]

P. B. Leonard and et al, 'Remote detection of small wetlands in the Atlantic coastal plain of North America: Local relief models, ground validation, and high-throughput computing', *Forest Ecology and Management*, vol. 284, pp. 107–115, doi: 10.1016/j.foreco.2012.07.034

[44]

E. R. Lundblad and et al, 'A Benthic Terrain Classification Scheme for American Samoa', *Marine Geodesy*, vol. 29, no. 2, pp. 89–111, 2006, doi: 10.1080/01490410600738021

[45]

S. N. Miller and et al, 'The Automated Geospatial Watershed Assessment tool', *Environmental Modelling & Software*, vol. 22, no. 3, pp. 365–377, doi: 10.1016/j.envsoft.2005.12.004

[46]

A. T. Naito and D. M. Cairns, 'Relationships between Arctic shrub dynamics and topographically derived hydrologic characteristics', *Environmental Research Letters*, vol. 6, no. 4, doi: 10.1088/1748-9326/6/4/045506

[47]

M. G. Rozo, A. C. R. Nogueira, and W. Truckenbrodt, 'The anastomosing pattern and the extensively distributed scroll bars in the middle Amazon River', *Earth Surface Processes and Landforms*, vol. 37, no. 14, pp. 1471–1488, doi: 10.1002/esp.3249

[48]

B. Rumsby, 'Vertical accretion rates in fluvial systems: -a comparison of volumetric and depth-based estimates', *Earth Surface Processes and Landforms*, vol. 25, no. 6, pp. 617–631, doi: 10.1002/1096-9837(200006)25:6<617::AID-ESP99>3.0.CO;2-Z

[49]

B. T. Rumsby and M. G. Macklin, 'Channel and floodplain response to recent abrupt climate change: The tyne basin, Northern England', *Earth Surface Processes and Landforms*, vol. 19, no. 6, pp. 499–515, doi: 10.1002/esp.3290190603

[50]

R. Sørensen, U. Zinko, and J. Seibert, 'On the calculation of the topographic wetness index: evaluation of different methods based on field observations', *Hydrology and Earth System Sciences*, vol. 10, no. 1, pp. 101–112, doi: 10.5194/hess-10-101-2006

[51]

D. G. Tarboton, R. L. Bras, and I. Rodriguez-Iturbe, 'On the extraction of channel networks from digital elevation data', *Hydrological Processes*, vol. 5, no. 1, pp. 81–100, doi: 10.1002/hyp.3360050107

[52]

J. Teng, J. Vaze, N. K. Tuteja, and J. C. Gallant, 'A GIS-Based Tool for Spatial and Distributed Hydrological Modelling: CLASS Spatial Analyst', *Transactions in GIS*, vol. 12, no. 2, pp. 209–225, doi: 10.1111/j.1467-9671.2008.01096.x

[53]

M. F. J. Wilson and et al, 'Multiscale Terrain Analysis of Multibeam Bathymetry Data for Habitat Mapping on the Continental Slope', *Marine Geodesy*, vol. 30, no. 1–2, pp. 3–35, doi: 10.1080/01490410701295962

[54]

F. Chaaban and et al, 'Using ArcGIS Modelbuilder and Aerial Photographs to Measure Coastline Retreat and Advance: North of France', *Journal of Coastal Research*, vol. 28, no. 6, pp. 1567–1579, 2012, Available: <http://www.bioone.org/doi/abs/10.2112/JCOASTRES-D-11-00054.1>

[55]

S. T. Hall and C. J. Post, 'Advanced GIS Exercise: Estimating Beach and Dune Erosion in Coastal South Carolina', *Journal of Natural Resources and Life Sciences Education*, vol. 37, pp. 49–52, 2008.

[56]

D. J. Wright and W. D. Heyman, 'Introduction to the Special Issue: Marine and Coastal GIS for Geomorphology, Habitat Mapping, and Marine Reserves', *Marine Geodesy*, vol. 31, no. 4, pp. 223–230, doi: 10.1080/01490410802466306

[57]

S. E. Greco and R. E. Plant, 'Temporal mapping of riparian landscape change on the Sacramento river, miles 196–218, California, USA', *Landscape Research*, vol. 28, no. 4, pp. 405–426, doi: 10.1080/0142639032000150149

[58]

J. E. Estes, *Some applications of aerial infrared imagery*. Lawrence, KS: Allen Press Inc, 1966.

[59]

C. R. C. Sheppard and et al, 'Habitat mapping in the Caribbean for management and conservation: Use and assessment of aerial photography', *Aquatic Conservation: Marine and Freshwater Ecosystems*, vol. 5, no. 4, pp. 277-298, doi: 10.1002/aqc.3270050404

[60]

J. C. Taylor, T. R. Brewer, and A. C. Bird, 'Monitoring landscape change in the National Parks of England and Wales using aerial photo interpretation and GIS', *International Journal of Remote Sensing*, vol. 21, no. 13-14, pp. 2737-2752, doi: 10.1080/01431160050110269

[61]

R. Kadmon and R. Harari-Kremer, 'Studying Long-Term Vegetation Dynamics Using Digital Processing of Historical Aerial Photographs', *Remote Sensing of Environment*, vol. 68, no. 2, pp. 164-176, doi: 10.1016/S0034-4257(98)00109-6

[62]

J. Sutherland, 'Error analysis of Ordnance Survey map tidelines, UK', *Proceedings of the ICE - Maritime Engineering*, vol. 165, no. 4, pp. 189-197, doi: 10.1680/maen.2011.10